

Measure Information								Workgroup Information			Measure Selection Criteria						
#	Measure Name	NQF #	Steward	Category	Data Source	Description	Measure Status	Measure Origin	Unit of Analysis	Based on readily available data	Nationally-vetted (e.g. NQF endorsed)	Aligned with measures in other WA and national measure sets	Significant potential to positively impact health outcomes	Significant potential to reduce costs	Sufficient denominator size	If provider-focused, the provider can impact the outcome	Notes about the Measure/ Measure Complexities
116	Use of Appropriate Medications for Asthma (ASM)	0036	NCQA	Asthma	Claims	Percentage of patients 5-64 years of age who were identified as having persistent asthma and were appropriately prescribed medication during the measurement period.	Yes	Aligned									
71	Medication Management for People with Asthma (MMA)	1799	NCQA	Asthma	Claims	The percentage of patients 5-64 years of age during the measurement year who were identified as having persistent asthma and were dispensed appropriate medications that they remained on during the treatment period. Two rates are reported. 1. The percentage of patients who remained on an asthma controller medication for at least 50% of their treatment period. 2. The percentage of patients who remained on an asthma controller medication for at least 75% of their treatment period.	No- Parking Lot	Aligned									
172	Asthma: Pharmacologic Therapy	0047	AMA-PCPI	Asthma	Claims and Clinical Data	Percentage of patients aged 5 through 64 years with a diagnosis of persistent asthma who were prescribed long-term control medication Three rates are reported for this measure: 1. Patients prescribed inhaled corticosteroids (ICS) as their long term control medication 2. Patients prescribed other alternative long term control medications (non-ICS) 3. Total patients prescribed long-term control medication	No- Parking Lot	GSK request									
173	Suboptimal Asthma Control (SAC) and Absence of Controller Therapy (ACT)	0548	#N/A	Asthma	Claims and Clinical Data	Rate 1 (SAC): The percentage of patients aged 5-50 years as of the last day of the measurement year with persistent asthma who were dispensed more than 3 canisters of a short-acting beta2 agonist inhaler during the same 90-day period. Rate 2 (ACT): The percentage of patients aged 5-50 years as of the last day of the measurement year with persistent asthma who were dispensed more than 3 canisters of short acting beta2 agonist inhalers over a 90-day period and who did not receive controller therapy during the same 90-day period.	No- Parking Lot	GSK request									
174	Relative Resource Use for People with Asthma	1560	NCQA	Asthma	Claims	The risk-adjusted relative resource use by health plan members with asthma during the measurement year.	No- Parking Lot	GSK request									
171	Asthma Medication Ratio (AMR)	1800	NCQA	Asthma	Claims and Clinical Data	The percentage of patients 5–64 years of age who were identified as having persistent asthma and had a ratio of controller medications to total asthma medications of 0.50 or greater during the measurement year.	No- Parking Lot	GSK request									
187	Asthma: Assessment of Asthma Control	0001	AMA-PCPI	Asthma	Claims and Clinical Data	Percentage of patients who were evaluated during at least one office visit for the frequency (numeric) of daytime and nocturnal asthma symptoms	No- Parking Lot	Measure Library									

Measure Information								Workgroup Information			Measure Selection Criteria						
#	Measure Name	NQF #	Steward	Category	Data Source	Description	Measure Status	Measure Origin	Unit of Analysis	Based on readily available data	Nationally-vetted (e.g. NQF endorsed)	Aligned with measures in other WA and national measure sets	Significant potential to positively impact health outcomes	Significant potential to reduce costs	Sufficient denominator size	If provider-focused, the provider can impact the outcome	Notes about the Measure/ Measure Complexities
119	Use of Spirometry Testing in the Assessment and Diagnosis of COPD (SPR)	0577	NCQA	COPD	Claims	The percentage of patients 40 years of age and older with a new diagnosis of COPD or newly active COPD, who received appropriate spirometry testing to confirm the diagnosis.	Yes	Aligned									
13	Anti-depressant Medication Management (AMM)	0105	NCQA	Depression	Claims	Percentage of patients 18 years of age and older who were diagnosed with major depression and treated with antidepressant medication, and who remained on antidepressant medication treatment. Two rates are reported. A. Percentage of patients who remained on an antidepressant medication for at least 84 days (12 weeks). b. Percentage of patients who remained on an antidepressant medication for at least 180 days (6 months).	Yes	Aligned									

Measure Information								Workgroup Information			Measure Selection Criteria						
#	Measure Name	NQF #	Steward	Category	Data Source	Description	Measure Status	Measure Origin	Unit of Analysis	Based on readily available data	Nationally-vetted (e.g. NQF endorsed)	Aligned with measures in other WA and national measure sets	Significant potential to positively impact health outcomes	Significant potential to reduce costs	Sufficient denominator size	If provider-focused, the provider can impact the outcome	Notes about the Measure/ Measure Complexities
257	Substance Use Disorders: Screening for Depression Among Patients with Substance Abuse or Dependence	NA	CMS	Depression	Claims or Registry	Percentage of patients aged 18 years and older with a diagnosis of current substance abuse or dependence who were screened for depression within the 12-month reporting period	Not yet considered	New addition per workgroup request									
259	Adult depression in primary care: percentage of patients with cardiovascular disease with documentation of screening for major depression or persistent depressive disorder using PHQ-2.	NA	Institute for Clinical Systems Improvement	Depression	Electronic health/medical record	This measure is used to assess the percentage of patients 18 years and older with cardiovascular disease with documentation of screening for major depression or persistent depressive disorder using Patient Health Questionnaire-2 (PHQ-2).	Not yet considered	New addition per workgroup request									
260	Adult depression in primary care: percentage of patients with type 2 diabetes with documentation of screening for major depression or persistent depressive disorder using PHQ-2.	NA	Institute for Clinical Systems Improvement	Depression	Electronic health/medical record	This measure is used to assess the percentage of patients 18 years and older with type 2 diabetes with documentation of screening for major depression or persistent depressive disorder using Patient Health Questionnaire-2 (PHQ-2).	Not yet considered	New addition per workgroup request									
31	Comprehensive Diabetes Care: Eye Exam	0055	NCQA	Diabetes	Claims	Percentage of patients 18-75 years of age with diabetes who had a retinal or dilated eye exam by an eye care professional during the measurement period or a negative retinal exam (no evidence of retinopathy) in the 12 months prior to the measurement period	Yes	Aligned									
37	Comprehensive Diabetes Care: Medical Attention for Nephropathy	0062	NCQA	Diabetes	Claims	The percentage of patients 18-75 years of age with diabetes who had a nephropathy screening test or evidence of nephropathy during the measurement period.	Yes	Aligned									
33	Comprehensive Diabetes Care: Hemoglobin A1c (HbA1c) Poor Control (>9.0%)	0059	NCQA	Diabetes	Claims and Clinical Data	Percentage of patients 18-75 years of age with diabetes who had hemoglobin A1c > 9.0% during the measurement period.	No- Parking Lot	Aligned									
30	Comprehensive Diabetes Care: Blood Pressure Control (<140/90 mm Hg)	0061	NCQA	Diabetes	Claims and Clinical Data	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) whose most recent blood pressure (BP) reading is <140/90 mm Hg during the measurement year.	No- Parking Lot	Aligned									

Measure Information								Workgroup Information			Measure Selection Criteria						
#	Measure Name	NQF #	Steward	Category	Data Source	Description	Measure Status	Measure Origin	Unit of Analysis	Based on readily available data	Nationally-vetted (e.g. NQF endorsed)	Aligned with measures in other WA and national measure sets	Significant potential to positively impact health outcomes	Significant potential to reduce costs	Sufficient denominator size	If provider-focused, the provider can impact the outcome	Notes about the Measure/ Measure Complexities
41	DM: Use of Ace Inhibitors/ARBs	NA		Diabetes	Claims and Clinical Data	Percent of individuals 40-80 years old with type 1 or type 2 diabetes who had at least one prescription for an ACE inhibitor or angiotensin receptor blocker (ARB) in the last 6 months	No- Parking Lot	Health Plan									
42	DM: Use of Statins	NA		Diabetes	Claims and Clinical Data	Percent of individuals 40-80 years old with type 1 or type 2 diabetes who had at least one prescription for a statin in the last 6 months.	No- Parking Lot	Health Plan									
34	Comprehensive Diabetes Care: Hemoglobin A1c testing	0057	NCQA	Diabetes:A1C	Claims	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) who received an HbA1c test during the measurement year.	Yes	Aligned									
65	Initiation and Engagement of Alcohol and Other Drug Dependence Treatment (IET)	0004	NCQA	Drug and Alcohol Use	Claims	The percentage of adolescent and adult patients with a new episode of alcohol or other drug (AOD) dependence who received the following. - Initiation of AOD Treatment. The percentage of patients who initiate treatment through an inpatient AOD admission, outpatient visit, intensive outpatient encounter or partial hospitalization within 14 days of the diagnosis. - Engagement of AOD Treatment. The percentage of patients who initiated treatment and who had two or more additional services with a diagnosis of AOD within 30 days of the initiation visit	Maybe	Health Plan									
129	SBIRT Service Penetration	NA	Washington State Defined	Drug and Alcohol Use	NA	The percentage of members who had an outpatient visit and who received a Screening, Brief Intervention or Referral to Treatment (SBIRT) service during the measurement year or the year prior to the measurement year	Not yet considered	5732									
166	Preventive Care & Screening: Unhealthy Alcohol Use: Screening & Brief Counseling	2152	AMA-PCPI	Drug and Alcohol Use	Patient Reported Data/Survey	Percentage of patients aged 18 years and older who were screened for unhealthy alcohol use at least once during the two year measurement period using a systematic screening method AND who received brief counseling if identified as an unhealthy alcohol user.	Not yet considered	Member Request: Ian Corbridge									
258	SUB-3 Alcohol & Other Drug Use Disorder Treatment Provided or Offered at Discharge and SUB-3a Alcohol & Other Drug Use Disorder Treatment at Discharge	1664	The Joint Commission	Drug and Alcohol Use	Clinical data	This measure is used to assess the percent of hospitalized patients 18 years of age and older who are identified with an alcohol or drug use disorder who received a prescription at discharge for Food and Drug Administration (FDA)-approved medication for treatment of alcohol or drug use disorder OR a referral for addictions treatment.	Not yet considered	New addition per workgroup request									
256	Substance Use Disorders: Counseling Regarding Psychosocial and Pharmacologic Treatment Options for Alcohol Dependence	NA	CMS	Drug and Alcohol Use	Claims or Registry	Percentage of patients aged 18 years and older with a diagnosis of current alcohol dependence who were counseled regarding psychosocial AND pharmacologic treatment options for alcohol dependence within the 12-month reporting period	Not yet considered	New addition per workgroup request									
18	ASCVD: Use of Statins	NA	American College of Cardiology and the American Heart Association	Hypertension and Cardiovascular Disease	Claims	Cholesterol-Lowering Medication for Patients with Coronary Artery Disease	Maybe	Aligned									

Measure Information								Workgroup Information			Measure Selection Criteria						
#	Measure Name	NQF #	Steward	Category	Data Source	Description	Measure Status	Measure Origin	Unit of Analysis	Based on readily available data	Nationally-vetted (e.g. NQF endorsed)	Aligned with measures in other WA and national measure sets	Significant potential to positively impact health outcomes	Significant potential to reduce costs	Sufficient denominator size	If provider-focused, the provider can impact the outcome	Notes about the Measure/ Measure Complexities
27	Cholesterol Management for Patients with Cardiovascular Conditions (LDL-C Screening)) (CMC)	NA	NCQA	Hypertension and Cardiovascular Disease	Claims and Clinical Data	The percentage of members 18–75 years of age who were discharged alive for AMI, coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) in the year prior to the measurement year, or who had a diagnosis of ischemic vascular disease (IVD) during the measurement year and the year prior to the measurement year, who had each of the following during the measurement year: <ul style="list-style-type: none">• LDL-C screening.	Maybe	Aligned									
126	Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia	1933	NCQA	Hypertension and Cardiovascular Disease	Claims	The percentage of members 18–64 years of age with schizophrenia and cardiovascular disease, who had an LDL-C test during the measurement year.	Not yet considered	5732									
145	Ischemic Vascular Disease (IVD): Use of Aspirin or Another Antithrombotic	0068	NCQA	Hypertension and Cardiovascular Disease	Clinical Data	Percentage of patients 18 years of age and older who were discharged alive for acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) in the 12 months prior to the measurement period, or who had an active diagnosis of ischemic vascular disease (IVD) during the measurement period, and who had documentation of use of aspirin or another antithrombotic during the measurement period.	Not yet considered	Aligned									
69	Ischemic Vascular Disease (IVD): Complete Lipid Panel and LDL Control	0075	NCQA	Hypertension and Cardiovascular Disease	Claims and Clinical Data	Percentage of patients 18 years of age and older who were discharged alive for acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) in the 12 months prior to the measurement period, or who had an active diagnosis of ischemic vascular disease (IVD) during the measurement period, and who had a complete lipid profile performed during the measurement period and whose LDL-C was adequately controlled (< 100 mg/dL).	Not yet considered	Aligned									
55	Heart Failure (HF): Angiotensin-Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) Therapy for Left Ventricular Systolic Dysfunction (LVSD)	0081	AMA-PCPI	Hypertension and Cardiovascular Disease	Claims and Clinical Data	Percentage of patients aged 18 years and older with a diagnosis of heart failure (HF) with a current or prior left ventricular ejection fraction (LVEF) < 40% who were prescribed ACE inhibitor or ARB therapy either within a 12 month period when seen in the outpatient setting OR at each hospital discharge	Not yet considered	Health Plan									
1	ACE-I/ARB: Persistent use with lab monitoring	NA	NCQA	Hypertension and Cardiovascular Disease	Claims		Not yet considered	Health Plan									
17	ASCVD: Use of Ace Inhibitors/ ARBs	NA		Hypertension and Cardiovascular Disease	Claims	Percent of individuals 18-80 years old who were diagnosed with atherosclerotic cardiovascular disease (ASCVD) who had at least one prescription for an ACE inhibitor or angiotensin receptor blocker (ARB) in the last 6 months	Not yet considered	Health Plan									

Measure Information								Workgroup Information			Measure Selection Criteria						
#	Measure Name	NQF #	Steward	Category	Data Source	Description	Measure Status	Measure Origin	Unit of Analysis	Based on readily available data	Nationally-vetted (e.g. NQF endorsed)	Aligned with measures in other WA and national measure sets	Significant potential to positively impact health outcomes	Significant potential to reduce costs	Sufficient denominator size	If provider-focused, the provider can impact the outcome	Notes about the Measure/ Measure Complexities
197	Hypertension (HTN): Blood Pressure Measurement	0013	AMA-PCPI	Hypertension and Cardiovascular Disease	Clinical Data	Percentage of patients aged 18 years and older with a diagnosis of hypertension with a blood pressure <140/90 mm Hg OR patients with a blood pressure >= 140/90 mm Hg and prescribed 2 or more anti-hypertensive medications during the most recent office visit within a 12 month period	Not yet considered	Measure Library									
193	Chronic Stable Coronary Artery Disease: ACE Inhibitor or ARB Therapy--Diabetes or Left Ventricular Systolic Dysfunction (LVEF <40%)	0066	American College of Cardiology	Hypertension and Cardiovascular Disease	Claims and Clinical Data	Percentage of patients aged 18 years and older with a diagnosis of CAD seen within a 12 month period who also have diabetes or a current or prior LVEF <40% who were prescribed ACE inhibitor or ARB therapy	Not yet considered	Measure Library									
189	Coronary Artery Disease (CAD): Beta-Blocker Therapy—Prior Myocardial Infarction (MI) or Left Ventricular Systolic Dysfunction (LVEF <40%)	0070	American College of Cardiology	Hypertension and Cardiovascular Disease	Clinical data	Percentage of patients aged 18 years and older with a diagnosis of coronary artery disease seen within a 12 month period who also have a prior MI or a current or prior LVEF <40% who were prescribed beta-blocker therapy	Not yet considered	Measure Library									
190	Ischemic Vascular Disease (IVD): Blood Pressure Management Control	0073	NCQA	Hypertension and Cardiovascular Disease	Claims and Clinical Data	The percentage of patients 18 to 75 years of age who were discharged alive with acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) during the 12 months prior to the measurement year, or who had a diagnosis of ischemic vascular disease (IVD) during the measurement year and the year prior to the measurement year and who had the following during the measurement year: - Blood pressure control (BP): reported as under control <140/90 mm Hg.	Not yet considered	Measure Library									
191	Chronic Stable Coronary Artery Disease: Lipid Control	0074	American College of Cardiology	Hypertension and Cardiovascular Disease	Claims and Clinical Data	Percentage of patients aged 18 years and older with a diagnosis of CAD seen within a 12 month period who have a LDL-C result <100 mg/dL OR patients who have a LDL-C result >=100 mg/dL and have a documented plan of care to achieve LDL-C <100mg/dL, including at a minimum the prescription of a statin	Not yet considered	Measure Library									
192	Heart Failure: Warfarin Therapy for Patients with Atrial Fibrillation	0084	AMA-PCPI	Hypertension and Cardiovascular Disease	Claims	Percentage of patients with HF who also have paroxysmal or chronic atrial fibrillation who were prescribed warfarin therapy.	Not yet considered	Measure Library									
139	Blood Pressure Screening by 13 years of age	1552	NCQA	Hypertension and Cardiovascular Disease	Claims and Clinical Data	The percentage of adolescents 13 years of age who had a blood pressure screening with results during the measurement year or the year prior to the measurement year	Not yet considered	Measure Library									
151	Taking medicine for high blood pressure control among adults aged >= 18	NA	Behavioral Risk Factor Surveillance System (BRFSS)	Hypertension and Cardiovascular Disease	Survey	Taking medicine for high blood pressure control among adults aged >= 18	Not yet considered	Member Request: Kristen Wendorf									
38	Controlling High Blood Pressure	0018	NCQA	Hypertension and Cardiovascular Disease	Clinical Data	The percentage of patients 18 to 85 years of age who had a diagnosis of hypertension (HTN) and whose blood pressure (BP) was adequately controlled (<140/90) during the measurement year.	No- Parking Lot	Aligned									

Measure Information								Workgroup Information			Measure Selection Criteria						
#	Measure Name	NQF #	Steward	Category	Data Source	Description	Measure Status	Measure Origin	Unit of Analysis	Based on readily available data	Nationally-vetted (e.g. NQF endorsed)	Aligned with measures in other WA and national measure sets	Significant potential to positively impact health outcomes	Significant potential to reduce costs	Sufficient denominator size	If provider-focused, the provider can impact the outcome	Notes about the Measure/ Measure Complexities
92	Pharmacy: Percent Generic (one rate for each: Antacid, Antidepressants, Statins, ACE and ARBS, ADHD)	NA	Washington Health Alliance Home Grown	Medication Management and Generic Use	Claims	Percentage of Generic Prescriptions for ACE inhibitors or angiotensin II receptor blockers (ARBs). Percentage of Generic Prescriptions for attention deficit hyperactivity disorder (ADHD) Medications Percentage of Generic Prescriptions for PPIs (proton pump inhibitors) Percentage of Generic Prescriptions for SSRIs, SNRIs, and other Second Generation Antidepressants Percentage of Generic Prescriptions for Statins	Yes	Aligned									
186	Proportion of Days Covered (PDC): 5 Rates by Therapeutic Category	0541	Pharmacy Quality Alliance	Medication Management and Generic Use	Clinical Data	Percentage of patients 18 years and older who met the proportion of days covered threshold of 80% during the measurement year. Rate is calculated separately for the following medication categories: Beta-Blockers, ACEI/ARB, Calcium-Channel Blockers, Diabetes Medication, Statins	Not yet considered	GSK request									
120	Warfarin: PT/INR monitoring	0612	Active Health Management	Medication Management and Generic Use	Claims	The percentage of patients taking warfarin who had PT/INR monitoring SPECIFIC EXCLUSIONS • Dialysis in the past 4 months • Hospitalization in the past 4 months • Phlebotomy in the past 4 months • Office Visits in the past 4 months	Not yet considered	Health Plan									
70	Lab Monitoring for Chronic Meds	NA		Medication Management and Generic Use	Claims		Not yet considered	Health Plan									
93	Pharmacy: Percent Mail-Order	NA		Medication Management and Generic Use	Claims		Not yet considered	Health Plan									
50	Follow-Up Care for Children Prescribed Attention-Deficit/Hyperactivity Disorder Medication (ADD)	0108	NCQA	Medication Management and Generic Use	Claims	Percentage of children newly prescribed ADHD medication that had at least three follow-up care visits within a 10-month period, one of which was within 30 days from the time the first ADHD medication was dispensed, including two rates: one for the initiation phase and one for the continuation and maintenance phase	No- Parking Lot	Aligned									
39	Disease Modifying Anti-Rheumatic Drug (DMARD) Therapy for Rheumatoid Arthritis (ART)	0054	NCQA	Other	Claims	The percentage of patients 18 years and older by the end of the measurement period, diagnosed with rheumatoid arthritis and who had at least one ambulatory prescription for a disease-modifying anti-rheumatic drug (DMARD).	Not yet considered	Health Plan									
74	Migraine: Frequent use of meds/receiving prophylactic meds	NA	Optum	Other	Claims	This measure calculates the percentage of members age 18 or older with migraines frequently taking acute (abortive) medications and taking a prophylactic medication for migraine control	Not yet considered	Health Plan									
113	Total Cost of Care Budgeted PMPM	NA		Other	Claims	Total Cost of Care Budgeted PMPM	Not yet considered	Health Plan									
202	Primary Open Angle Glaucoma (POAG): Optic Nerve Evaluation	0086	AMA-PCPI	Other	Claims and Clinical Data	Percentage of patients aged 18 years and older with a diagnosis of POAG who have an optic nerve head evaluation during one or more office visits within 12 months	Not yet considered	Measure Library									

Measure Information								Workgroup Information			Measure Selection Criteria						
#	Measure Name	NQF #	Steward	Category	Data Source	Description	Measure Status	Measure Origin	Unit of Analysis	Based on readily available data	Nationally-vetted (e.g. NQF endorsed)	Aligned with measures in other WA and national measure sets	Significant potential to positively impact health outcomes	Significant potential to reduce costs	Sufficient denominator size	If provider-focused, the provider can impact the outcome	Notes about the Measure/ Measure Complexities
203	CAHPS® In-Center Hemodialysis Survey	0258	CMS	Other	survey	Percentage of patient responses to multiple testing tools. Tools include the In-Center Hemomdialysis Composite Score: The proportion of respondents answering each of response options for each of the items summed across the items within a composite to yield the composite measure score. (Nephrologists' Communication and Caring, Quality of Dialysis Center Care and Operations, Providing Information to Patients) Overall Rating: a summation of responses to the rating items grouped into 3 levels	Not yet considered	Measure Library									
220	CAHPS® Nursing Home Survey: Discharged Resident Instrument	0691	AHRQ	Other	Survey	The CAHPS® Nursing Home Survey: Discharged Resident Instrument is a mail survey instrument to gather information on the experience of short stay (5 to 100 days) residents recently discharged from nursing homes. This survey can be used in conjunction with the CAHPS Nursing Home Survey: Family Member Instrument and the Long Stay Resident Instrument. The survey instrument provides nursing home level scores on 4 global items. In addition, the survey provides nursing home level scores on summary measures valued by consumers; these summary measures or composites are currently being analyzed. The composites may include those valued by long stay residents: (1) Environment; (2) Care; (3) Communication & Respect; (4) Autonomy and (5) Activities.	Not yet considered	Measure Library									
228	HIV Viral Load Suppression	2082	HRSA - HIV/AIDS Bureau	Other	Claims and Clinical Data	Percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.	Not yet considered	Measure Library									
224	CAHPS® Nursing Home Survey	0692-0693	AHRQ	Other	Survey	#N/A	Not yet considered	Measure Library									
170	Advance Care Plan	0326	NCQA	Other	Claims and Clinical Data	Percentage of patients aged 65 years and older who have an advance care plan or surrogate decision maker documented in the medical record or documentation in the medical record that an advance care plan was discussed but the patient did not wish or was not able to name a surrogate decision maker or provide an advance care plan.	Not yet considered	Member Request: Alice Lind									